Seniors Wellness Group of Michigan 221 S. Main Street, Suite 201 Royal Oak, MI 48067 248-396-6459

Risk Benefit Analysis (RBA)

(2013)

| Patient: | Facility: | |
|---|--|--|
| Examiner: | Date: | Room: |
| | | |
| O. L 100 7 . 1 106 15(A) | | |
| Ordered/Prescribed Medication(s): | | |
| Common or Serious Adverse Side Effects and Risks of | f Psychotropic Medications: | |
| □ AChE Inhibitors: GI upset, diarrhea, muscle cramps, anorea □ Anxiolytics: sedation, ataxia, paradoxical agitation, confusion motor skills, risk of falls | xia, orthostatic hypotension, syncop n, amnesia, addiction, respiratory s | pe, seizures, GI bleeding, arrhythmia uppression/aspiration, decreased |
| □ Antipsychotics: sedation, parkinsonism, akathisia, akinesia of urinary retention, metabolic syndrome, neuroleptic malignant suppression/aspiration; Black Box Warning: increased mor □ Hypnotics: daytime fatigue, cognitive impairment, falls and | t syndrome, stroke, cardiac arrest, p tality in elderly | d vision, dry mouth, constipation, riapism, respiratory |
| □ Namenda: dizziness, constipation, headaches, confusion | Hactures | |
| MAOIs: dry mouth, dizziness, headaches, sedation, nausea, h restriction | sypertensive crisis, orthostatic hypo | tension, seizures, tyramine |
| ☐ Tegretol: Dizziness, GI upset, nausea, vomiting, drowsiness, fatal) skin reactions (rare), aplastic anemia and other blood di | | on, tremors, very serious (possibly |
| □ Valproic Acid, Lamictal, Topamax: Hematological and live Black Box Warning: Valproic Acid – rare but serious live | er abnormalities, GI upset, sedatior | n, appetite changes, thrombocytopenia, |
| Lithium: GI upset, metallic taste, dizziness, increased appetit | e, fine tremor, polyuria, polydipsia | , kidney damage, hypothyroidism, visua |
| changes; Black Box Warning: toxicity □ SNRIs: nausea, sedation, dry mouth, diminished libido, anorg | gasmia, hypertension | : |
| SSRIs: headaches, fatigue, ejaculatory delay, impotence, anor | rgasmia, GI distress, abnormal blee | eding, SIADH (Citalopram: - dose- |
| dependent QT interval prolongation – maximum dosage 20 Tricyclics: dry mouth, blurred vision, constipation, sedation, | urinary retention, impotence, tachy | cardia, orthostatic hypotension |
| ☐ Trazodone: ataxia, orthostatic hypotension, drowsiness, priar ☐ Remeron: increased appetite, somnolence, hypo/hypertension | | |
| Wellbutrin: insomnia, agitation, headaches, orthostatic hypothesis | tension, hypertension, seizures, ras | h/allergic reaction (rare but serious) |
| ☐ Stimulants: insomnia, agitation, headaches, GI upset, anorexi | ia, hypertension, visual disturbance | es, irritability, seizures, blood dyscrasia, |
| Black Box Warning: addiction, psychosis Others: | · | |
| | | |
| Risk Benefit Statement and Decision to Treat (comorbidities, hypersensitivities, potential drug-drug | (as applicable, include consinteractions, etc.): | siderations of significant medic |
| | | |
| | | |
| <u> </u> | | |
| | | |
| | | |
| | | |
| · | | |
| | | |
| | <u> </u> | |
| | | |
| Description Committee to | Do4 | |
| Psychiatric Consultant: | Date: | |